

# Psychiatric Disorders and Learning Problems in Children and Adolescents with Sickle Cell Disease

Özalp Ekinci<sup>1</sup>, Tanju Çelik<sup>2</sup>

Mustafa Kemal Üniversitesi <sup>1</sup>Çocuk Psikiyatrisi Uzmanı, <sup>2</sup>Pediyatri Anabilim Dalı, Antakya

## ÖZET

*Orak Hücre Anemili çocuk ve ergenlerde psikiyatrik bozukluklar ve öğrenme problemleri*

Orak Hücre Anemisi (OHA) kronik bir süreç ve ağrılı krizleri de içeren ve hayatı olumsuz etkileyen komplikasyonlarla seyreden bir hastalıktır. OHA'lı çocuk ve ergenler uyum problemleri ve bozulmuş psikososyal işlevsellik açısından risk altında olarak görünmektedir. Mevcut literatür OHA tanılı çocuk ve ergenlerin normal kontrollere göre daha düşük hayat kalitesi ve daha yüksek sıklıkta nöro-psikiyatrik problemler gösterdiğine işaret etmektedir. Nöro-psikiyatrik problemler arasında depresyon, anksiyete bozuklukları, öğrenme bozuklukları, eksternalize davranış bozuklukları, enüresis ve dikkat güçlükleri sıklıkla bildirilmektedir. Bu yazı OHA'lı çocuk ve ergenlerde psikiyatrik bozukluklar ve öğrenme problemleri üzerine literatürü gözden geçirmeyi amaçlamaktadır.

**Anahtar kelimeler:** Orak Hücre Anemisi, çocuk, ergen, psikiyatrik bozukluklar

## ABSTRACT

*Psychiatric disorders and learning problems in children and adolescents with Sickle Cell Disease*

Sickle cell disease (SCD) is characterized by a chronic course and disabling complications including recurrent pain attacks. Children and adolescents with SCD appear to be at risk for adjustment problems and impaired psychosocial functioning. The available research indicates that children and adolescents with SCD have a lower quality of life and a higher frequency of neuro-psychiatric problems than that of normal controls. Among the neuro-psychiatric problems, depression, anxiety disorders, learning disorders, externalizing behavior problems, enuresis and attention difficulties are commonly reported. This article aims to review the current literature on psychiatric disorders and learning problems in children and adolescents with SCD.

**Key words:** Sickle Cell Disease, children, adolescents, psychiatric disorders

Bakırköy Tıp Dergisi 2012;8:95-100

## INTRODUCTION

Sickle cell disease (SCD) is a chronic, autosomal recessive disorder characterized by the predominance of the protein hemoglobin S (HbS) in red blood cells. SCD primarily affects people of African, Caribbean, Asian, Middle Eastern, and Mediterranean descent (1). SCD is characterized by a chronic course and disabling complications including an increased risk of infections. Recurrent pain crisis that requires emergency management may be considered as the hallmark of SCD (1,2). Children and adolescents with SCD face many challenges of living with a chronic condition that requires

lifelong medical management (2). As observed in children and adolescents with other chronic illnesses, the current literature suggests that children and adolescents with SCD may be at risk for adjustment problems and impaired psychosocial functioning (3,4).

The available research indicates that children and adolescents with SCD have a lower quality of life and a higher frequency of neuro-psychiatric problems when compared to the normal controls (2,5). Among the neuro-psychiatric problems, depression, anxiety disorders, learning problems, externalizing behavior problems, enuresis and attention difficulties are commonly reported (6-10). This article aims to review the current literature on psychiatric disorders and learning problems in children and adolescents with SCD.

## Quality of life

Health has been defined as being not only the

Yazışma adresi / Address reprint requests to: Özalp Ekinci, Kışlasaray Mahallesi, Hatay, Antakya

Telefon / Phone: +90-506-611-1500

Elektronik posta adresi / E-mail address: ozalpekinci@yahoo.com

Geliş tarihi / Date of receipt: 13 Aralık 2011 / December 13, 2011

Kabul tarihi / Date of acceptance: 15 Şubat 2012 / February 15, 2012

absence of disease and infirmity but also the presence of physical, mental, and social well-being (11). Health-related quality of life (HRQOL) refers to the physical, psychological, and social domains of health seen in areas influenced by a person's experiences, beliefs, expectations, and perceptions (12). Number of studies to date has examined HRQOL in chronic diseases of childhood and adolescence including obesity, cancer, and HIV infection (13,14,15). HRQOL in children and adolescents with SCD have also been studied in a handful of studies. The majority of these studies have reported a lower HRQOL when compared to the normal controls (16,17,18). In a more recent study, Dale et al reported that both children with SCD and their parents rated overall HRQOL and all subdomains of HRQOL lower than healthy children and their parents (2). There is also evidence that some differences may exist between the reports of children and parents on HRQOL. Panepinto et al. found that, compared with children, parents reported worse HRQOL in overall perception of health, physical functioning, behavior, and self-esteem domains (19).

### **Depression**

Majority of the available studies have shown that children and adolescents with SCD have a higher risk of depressive symptoms (6,7,20,21). These include both major depression and other forms of psychiatric diagnosis with depressive symptoms. In a recent study, approximately half of children and adolescents with SCD were diagnosed with either dysthymia (90%) or major depressive disorder (10%) (6). In that study, the age at onset of depressive disorders appears to be younger among patients with SCD when compared with the normal population. The mean age for dysthymic disorder and major depressive disorder were reported as 9 and 14 years of age, respectively (6). Like the children with other chronic illnesses, depression in children and adolescents with SCD is usually accompanied with impaired functioning on multiple domains. Adolescents with SCD have been reported to have significant problems in social relationships, isolation, and school failure, with 29% meeting the criteria for depression based on symptom scales (7).

Although the risk factors of depression in SCD are not completely known, there is evidence that disease related factors may have an influence. There appears to be a

positive correlation between the frequency of pain episodes and depressive symptoms (22). Jerrel et al. also has found that the children and adolescents with SCD with depression had a higher frequency of vaso-occlusive pain attacks and acute chest syndrome visits per year and developed more complications with related organ damage (6). As shown for other chronic medical conditions, parental psychiatric characteristics have been found to be associated with the emotional symptoms in children with SCD. Unal et al. has found a relation between anxiety, depression, and marital adjustment levels in mothers and depression in children (22).

### **Anxiety Disorders**

As observed in other chronic diseases, children and adolescents with SCD usually experience anxiety symptoms (7,20,21,22). The unpredictable painful nature of SCD, frequent hospitalizations, fear of worsening of the condition, worries about the future social life as an individual and the stigma associated with SCD appear to contribute to an increased risk of anxiety disorders (4,23). The predictors of anxiety symptoms may differ between childhood and adolescence. Younger children with SCD may experience separation anxiety symptoms, especially on separation from their mothers (24). This may partly be explained by the mothers' role in the health care of the children. Mothers' own worries about their child and overprotective parenting styles may also contribute to the separation anxiety of the child. Regarding to the adolescence period, body image and peer relations may be considered as the main domains of life. Since adolescents with SCD usually have a shorter stature than their peers, they may feel embarrassment in peer relations and, in turn, may withdraw from social activities (25). As the adolescent grow older, this anxiety on body image may disrupt their romantic relations. In addition to these psychosocial factors, disease related factors have also been suggested to be linked with anxiety symptoms in SCD. Mahdi et al. has found that the frequency of vasoocclusive attacks are related with an increased risk of anxiety symptoms (26).

### **Externalizing Behavior Problems**

Most of the studies generally suggest a higher prevalence of internalizing problems but no increased

risk of externalizing problems in children and adolescents with SCD (27). However, methodological issues of these studies, including the psychiatric scales used, might have limited the identification of externalizing problems in children with SCD. In an early study by Brown et al. children and adolescents with SCD have been found to have a higher frequency of externalizing behavioral problems than their healthy siblings rated by caregivers (28). In a more recent study, Hijman et al. has shown that the children with SCD had more severe externalizing problems than the normal population according to teacher rated data (8). The etiopathogenesis of externalizing problems in children with SCD has not been studied extensively. However, there is increasing evidence that externalizing behavioral problems, especially the symptoms associated with impulsivity, may be related with cerebral infarcts on the frontal lobes (29,30,31).

### **Enuresis**

Enuresis is commonly reported among children with SCA. The prevalence of enuresis is estimated between 29-50% in SCD which is much higher than the normal population data (32-35). Etiological explanations of the high rates of enuresis in SCD is largely unknown. However, some disease related factors have been suggested to be related with a higher frequency of enuresis in SCD. Babela et al. showed a positive correlation between a higher frequency pain attacks and enuresis (34). In contrast, a more recent study did not find a relation between pain attacks and enuresis (32). There is evidence that enuresis has a broad negative impact on children with SCD. Children with SCD who had enuresis have been reported to experience higher levels of total psychosocial problems (36).

### **Neurocognitive deficits and learning problems**

The available research has shown that the children and adolescents with SCD have a higher risk of learning problems and academic underachievement (37). Academic underachievement appears to reflect the interaction of neurocognitive deficits associated with SCD and the school attainment problems related with frequent hospitalizations (10,38,39,40). Neurocognitive deficits in SCD may be broadly classified as global

intellectual disabilities and specific neurocognitive impairments. Among the specific neurocognitive impairments, problems on attention, executive functions, memory, language and visuomotor functions have been studied in different studies (10,40-43).

One of the most devastating complications of SCD, which is directly related with neurocognitive impairments, is cerebral infarction. It is shown that cerebral infarcts are present on magnetic resonance imaging (MRI) scans in one third of children and adolescents with SCD by the age of 18 years (44-48). Both overt and silent infarcts evident on neuroimaging have been described in SCD. Overt infarct, cerebral infarction accompanied by neurological symptoms, is estimated to occur in 11% of children with SCD-SS (49). In contrast to the overt infarcts, silent infarcts have been defined as cerebral infarcts shown with MRI in the absence of clinical neurologic symptoms. Silent infarcts occur more commonly than overt infarcts, with a prevalence of 21.8% in children and adolescents with homozygous SCD (44).

Various studies have investigated intellectual function or general intelligence (IQ), in children with SCD. These studies mainly included the children above 6 years of age (10). However, a limited number of studies have also focused on younger children (47,50). One of the studies on younger children has reported delays in the developmental milestones (50). On the other hand, most of the studies on older children with SCD have reported varying degrees of impairment on IQ and subscales of IQ (51,52,53). In the light of the available research, it is suggested that impairment in intellectual development may appear in infancy and increase with age (10). IQ in children with SCD varies with the degree of neurological compromise. Children with evidence of clinical or silent infarcts were found to demonstrate lower total IQ scores and certain IQ subscores compared to normal population (40). There is also evidence that there may be differences on IQ between children with clinical and silent infarcts. Armstrong et al. have shown that, when compared with children with silent infarcts, the children with clinical infarcts demonstrated more deficits on certain IQ scores (43).

It is well known that executive functions are important for almost all of the cognitive functions. Majority of the studies to date have shown a higher frequency of impairments on executive functions in children with SCD when compared to the normal population (10). Among

the executive functions, working memory deficits appear to be more prominent (40). Brandling-Bennett et al. reported a decreased working memory span (54). White et al, more specifically, reported a decreased word length span (55). Deficits on working memory were also found to be related with cerebral injury and especially with frontal infarcts (54,55)

Attention difficulties are commonly reported in children and adolescents with SCD (8,10,56). Among attention problems, particularly sustained attention was suggested to be impaired in children SCD (10,29). Although not completely proven, deficits in attention are often related to the presence and severity of cerebral injury. Frontal lobe abnormalities, especially anterior infarcts, appear to underlie in the attention deficits (38,56,57). There is also evidence linking diffuse lesions to disruption of attentional processes on orienting tasks in SCD (58).

The available literature on memory functions in SCD is relatively limited. Schatz and Roberts have shown that children with SCD showed difficulties for digit span-backward (59). There is also evidence that deficits on visual-spatial memory are more frequently reported than deficits on verbal memory functions (41). As the other cognitive deficits, deficits on memory have been suggested to be related with neurological integrity including the presence and the severity of cerebral infarcts (10).

### **Management of psychiatric disorders in SCD**

Medical advances in the treatment of SCD have transformed it from a condition associated with very early morbidity and mortality into a life-long, chronic condition (7). However, the studies on the treatment of psychiatric disorders in SCD are limited. Moreover, the majority of children and adolescents with SCD do not reach psychiatric evaluation and treatment. It is estimated that only one-third of patients with SCD receive treatment from a mental health professional (27). Patient and family education is the hallmark of interventions. For the management of psychiatric problems, the utilization of psychological interventions including cognitive-behavioral therapy and medical treatment including psychotropic medications may be needed. Specific education services must be warranted for the children with intellectual disabilities. Childs' developmental needs and specific learning problems must be established

before the initiation of education. The children with cerebral infarcts, frequent hospitalizations and premorbid psychiatric symptoms must be regularly evaluated by a child and adolescent psychiatry specialist.

A limited number of studies investigated the safety and efficacy of psychotropic medications in children and adolescents with SCD. There appears to be a bidirectional relationship between emotional symptoms and chronic pain in SCD. It is suggested that children with SCD who experience chronic pain are usually depressed, and alleviating the depressive symptoms may have a salutary effect on the pain (6). There is evidence that the combination of antidepressants or anticonvulsants with opioids may be associated with fewer acute vaso-occlusive pain visits (60). Several antidepressants have been found to be effective in the management of disparate pain disorders, including amitriptyline, imipramine, venlafaxine, bupropion and duloxetine (61,62). However most of these studies included adult patients. The research on the selective serotonin reuptake inhibitors (SSRIs), which are considered as firstline medications for the treatment of depression and anxiety symptoms in the normal population of children and adolescents, is unfortunately limited in SCD. Jerrel et al. in their recent study did not reach a conclusive finding on the efficacy of SSRIs due to the high rates of non-compliance in their cohort (6).

When initiating a treatment with antidepressants, a careful evaluation by a child psychiatrist is needed to avoid the potential adverse reactions of the drug and the unwanted reactions with the other medications used. This evaluation includes an initial hematologic examination and, in needed cases, the consultation of a pediatric hematology specialist. Similar to the initial treatment, follow up of the patients may warrant timely hematologic examinations and consultation of a pediatric hematology specialist.

Similar to the treatment of other psychiatric disorders in children with SCD, the literature on the treatment of enuresis is limited. Primary caregivers use a wide range of interventions for nocturnal enuresis, however, there is a lack of the empirically supported treatments for enuresis (63). In an early study by Figueroa, intranasal desmopressin acetate has been shown to be effective in children and adolescents with SCD (35). Imipramine (a tricyclic antidepressant), has also been shown to be effective in reducing symptoms (63).

## CONCLUSION

Children and adolescents with SCD experience a high frequency of neuro-psychiatric problems. Neuro-psychiatric problems in SCD are usually underdiagnosed and undertreated, although they cause an additional burden on the quality of life of these patients. The

collaboration of different disciplines including general practitioners, pediatric hematologists and child or adolescent psychiatry specialists is crucial for the early diagnosis and effective treatment of these conditions. Future studies are needed to clarify the treatment options of psychiatric problems in children and adolescents with SCD.

## REFERENCES

1. Stuart MJ, Nagel RL. Sickle-cell disease. *Lancet* 2004; 364: 1343-1360.
2. Dale JC, Cochran CJ, Roy L, Jernigan E, Buchanan GR. Health-related quality of life in children and adolescents with sickle cell disease. *J Pediatr Health Care* 2011; 25: 208-215.
3. Thomas VJ, Taylor LM. The psychological experience of people with sickle cell disease and its impact on quality of life: Qualitative findings of a focus groups. *Br J Health Psychol* 2002; 7: 345-363.
4. Midence K, Fuggle P, Davies SC. Psychosocial aspects of sickle cell disease (SCD) in childhood and adolescence: a review. *Br J Clin Psychol* 1993; 32: 271-280.
5. Trzepacz A M, Vannatta K, Gerhardt CA, Ramey C, Noll RB. Emotional, social, and behavioral functioning of children with sickle cell disease and comparison peers. *J Pediatr Hematol Oncol* 2004; 26: 642-648.
6. Jerrell JM, Tripathi A, McIntyre RS. Prevalence and treatment of depression in children and adolescents with sickle cell disease: a retrospective cohort study. *Prim Care Companion CNS Disord* 2011; 13 pii: PCC.10m01063.
7. Benton TD, Ifeagwu JA, Smith-Whitley K. Anxiety and depression in children and adolescents with sickle cell disease. *Curr Psychiatry Rep* 2007; 9: 114-121.
8. Hijmans CT, Grootenhuis MA, Oosterlaan J, et al. Behavioral and emotional problems in children with sickle cell disease and healthy siblings: Multiple informants, multiple measures. *Pediatr Blood Cancer* 2009; 53: 1277-1283.
9. Field JJ, Austin PF, An P, Yan Y, DeBaun MR. Enuresis is a common and persistent problem among children and young adults with sickle cell anemia. *Urology* 2008; 72: 81-84.
10. Berkelhammer LD, Williamson AL, Sanford SD, et al. Neurocognitive sequelae of pediatric sickle cell disease: a review of the literature. *Child Neuropsychol* 2007; 13: 120-131.
11. World Health Organization. Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference, New York, 19-22 June, 1946; signed on 22 July 1946 by the representatives of 61 States (Official Records of the World Health Organization). Geneva, Switzerland: World Health Organization.
12. Testa MA, Simonson DC. Assessment of quality of life outcomes. *N Engl J Med*; 334: 835-840.
13. Williams J, Wake M, Hesketh K, Maher E, Waters E. Health-related quality of life of overweight and obese children. *JAMA* 2005; 293: 70-76.
14. Speechley KN, Barrera M, Shaw AK, et al. Health-related quality of life among children and adolescent survivors of childhood cancer. *J Clin Oncol* 2006; 24: 2536-2543.
15. Lee GM, Gortmaker SL, McIntosh K, Hughes MD, Oleske JM. Quality of life for children and adolescents: Impact of HIV infection and antiviral treatment. *Pediatrics* 2006; 117: 273-283.
16. Wethers D. Sickle cell disease in childhood: Part II. Diagnosis and treatment of major complications and recent advances in treatment. *Am Fam Physician* 2000; 62: 1309-1314.
17. Ramsey LT, Woods KF, Callahan LA, Mensah GA, Barbeau P, Gutin B. Quality of life improvement for patients with sickle cell disease. *Am J Hematol* 2001; 66: 155-156.
18. Palermo TM, Schwartz L, Drotar D, McGowan K. Parental report of health-related quality of life in children with sickle cell disease. *J Behav Med* 2002; 25: 269-283.
19. Panepinto JA, O'Mahar KM, DeBaun MR, Rennie KM, Scott JP. Validity of the child health questionnaire for use in children with sickle cell disease. *J Pediatr Hematol Oncol* 2004; 26: 574-578.
20. Benton TD, Boyd R, Ifeagwu J, Feldtmose E, Smith-Whitley K. Psychiatric diagnosis in adolescents with sickle cell disease: a preliminary report. *Curr Psychiatry Rep* 2011; 13: 111-115.
21. Alao AO, Dewan MJ, Jindal S, Effron M. Psychopathology in sickle cell disease. *West Afr J Med* 2003; 22: 334-337.
22. Unal S, Toros F, Kütük MÖ, Uyaniker MG. Evaluation of the psychological problems in children with sickle cell anemia and their families. *Pediatr Hematol Oncol* 2011; 28: 321-328.
23. Serjeant GR. The clinical features of sickle cell disease. *Baillieres Clin Hematol* 1993; 6: 93-115.
24. Thompson RJ Jr, Gil KM, Burbach DJ, Keith BR, Kinney TR. Role of child and maternal processes in the psychological adjustment of children with sickle cell disease. *J Consult Clin Psychol* 1993; 61: 468-474.
25. Hurtig AL, Park KB. Adjustment and coping in adolescents with sickle cell disease. *Ann N Y Acad Sci* 1989; 565: 172-182.
26. Mahdi N, Al-Ola K, Khalek NA, Almawi WY. Depression, anxiety, and stress comorbidities in sickle cell anemia patients with vaso-occlusive crisis. *J Pediatr Hematol Oncol* 2010; 32: 345-349.
27. Barakat L, Lash LA, Lutz MJ, et al. Psychosocial adaptation of children and adolescents with sickle cell disease. In: Brown RT (Ed). *Comprehensive handbook of childhood cancer and sickle cell disease*. New York: Oxford University Press; 2006: P.471-495.
28. Brown RT, Kaslow NJ, Doepke K, et al. Psychosocial and family functioning in children with sickle cell syndrome and their mothers. *J Am Acad Child Adolesc Psychiatry* 1993; 32: 545-553.
29. DeBaun MR, Schatz J, Siegel MJ, et al. Cognitive screening examinations for silent cerebral infarcts in sickle cell disease. *Neurology* 1998; 50: 1678-1682.

30. Noll RB, Stith L, Gartstein MA, et al. Neuropsychological functioning of youths with sickle cell disease: Comparison with non-chronically ill peers. *J Pediatr Psychol* 2001; 26: 69-78.
31. Schatz J, Craft S, Koby M, et al. Neuropsychologic deficits in children with sickle cell disease and cerebral infarction: Role of lesion site and volume. *Child Neuropsychol* 1999; 5: 92-103.
32. Field JJ, Austin PF, An P, Yan Y, DeBaun MR. Enuresis is a common and persistent problem among children and young adults with sickle cell anemia. *Urology* 2008; 72: 81-84.
33. Akinyanju O, Agbato O, Ogunmekan AO, Okoye JU. Enuresis in sickle cell disease. I. Prevalence studies. *J Trop Pediatr* 1989; 35: 24-26.
34. Mabilia Babela JR, Loumingou R, Pemba-Loufoua A, Londjongo W, Nzingoula S, Senga P. Enuresis in children with sickle cell disease. *Arch Pediatr* 2004; 11: 1168-72.
35. Figueroa TE, Benaim E, Griggs ST, Hvizdala EV. Enuresis in sickle cell disease. *J Urol* 1995; 153: 1987-1989.
36. Jordan SS, Hilker KA, Stoppelbein L, Elkin TD, Applegate H, Iyer R. Nocturnal enuresis and psychosocial problems in pediatric sickle cell disease and sibling controls. *J Dev Behav Pediatr* 2005; 26: 404-411.
37. Schatz J, Finke R, Roberts CW. Interactions of biomedical and environmental risk factors for cognitive development: A preliminary study of sickle cell disease. *J Dev Behav Pediatr* 2004; 25: 303-310.
38. Schatz J, Brown RT, Pascual JM, Hsu L, DeBaun MR. Poor school and cognitive functioning with silent cerebral infarcts and sickle cell disease. *Neurology* 2001; 56: 1109-1111.
39. Shapiro BS, Dinges DF, Orne EC, et al. Home management of sickle cell-related pain in children and adolescents: Natural history and impact on school attendance. *Pain* 1995; 61: 139-144.
40. Wang W, Enos L, Gallagher D, et al. Neuropsychologic performance in school-age children with sickle cell disease: a report from the Cooperative Study of Sickle Cell Disease. *J Pediatr* 2001; 139: 391-397.
41. Watkins KE, Hewes DK, Connelly A, et al. Cognitive deficits associated with frontal-lobe infarction in children with sickle cell disease. *Dev Med Child Neurol* 1998; 40: 536-543.
42. Hariman LM, Griffith ER, Hurtig AL, Keehn MT. Functional outcomes of children with sickle-cell disease affected by stroke. *Arch Physic Med Rehabil* 1991; 72: 498-502.
43. Armstrong FD, Thompson RJ Jr, Wang W, et al. Cognitive functioning and brain magnetic resonance imaging in children with sickle cell disease. Neuropsychology Committee of the Cooperative Study of Sickle Cell Disease. *Pediatrics* 1996; 97: 864-870.
44. Pegelow CH, Macklin EA, Moser FG, et al. Longitudinal changes in brain magnetic resonance imaging findings in children with sickle cell disease. *Blood* 2002; 99: 3014-3018.
45. Seibert JJ, Miller SF, Kirby RS, et al. Cerebrovascular disease in symptomatic and asymptomatic patients with sickle cell anemia: Screening with duplex transcranial Doppler US- Correlation with MR imaging and MR angiography. *Radiology* 1993; 189: 457-466.
46. Steen RG, Emudianughe T, Hankins GM, et al. Brain imaging findings in pediatric patients with sickle cell disease. *Radiology* 2003; 228: 216-225.
47. Wang WC, Langston JW, Steen RG, et al. Abnormalities of the central nervous system in very young children with sickle cell anemia. *J Pediatr* 1998; 132: 994-998.
48. Wang WC, Gallagher DM, Pegelow CH, et al. Multicenter comparison of magnetic resonance imaging and transcranial Doppler ultrasonography in the evaluation of the central nervous system in children with sickle cell disease. *J Pediatr Hematol Oncol* 2000; 22: 335-339.
49. Ohene-Frempong K, Weiner SJ, Sleeper LA, et al. Cerebrovascular accidents in sickle cell disease: rates and risk factors. *Blood* 1998; 91: 288-294.
50. Hogan AM, Kirkham FJ, Prenalger M, et al. An exploratory study of physiological correlates of neurodevelopmental delay in infants with sickle cell anaemia. *Br J Haematol* 2006; 132: 99-107.
51. Kral MC, Brown RT, Hynd GW. Neuropsychological aspects of pediatric sickle cell disease. *Neuropsychol Rev* 2001; 11: 179-196.
52. Kral MC, Brown RT, Nietert PJ, et al. Transcranial doppler ultrasonography and neurocognitive functioning in children with sickle cell disease. *Pediatrics* 2003; 112: 324-331.
53. Steen RG, Fineberg-Buchner C, Hankins G, Weiss L, Prifitera A, Mulhern RK. Cognitive deficits in children with sickle cell disease. *J Child Neurol* 2005; 20: 102-107.
54. Brandling-Bennett EM, White DA, Armstrong MM, Christ SE, DeBaun M. Patterns of verbal long-term and working memory performance reveal deficits in strategic processing in children with frontal infarcts related to sickle cell disease. *Dev Neuropsychol* 2003; 24: 423-434.
55. White DA, Salorio CF, Schatz J, DeBaun M. Preliminary study of working memory in children with stroke related to sickle cell disease. *J Clin Exp Neuropsychol* 2000; 22: 257-264.
56. Nabors NA, Freymuth AK. Attention deficits in children with sickle cell disease. *Percept Mot Skills*. 2002; 95: 57-67.
57. Brown RT, Davis PC, Lambert R, et al. Neurocognitive functioning and magnetic resonance imaging in children with sickle cell disease. *J Pediatr Psychol* 2000; 25: 503-513.
58. Craft S, Schatz J, Glauser TA, Lee B, DeBaun MR. The effects of bifrontal stroke during childhood on visual attention: Evidence from children with sickle cell anemia. *Dev Neuropsychol* 1994; 10: 285-297.
59. Schatz J, Roberts CW. Short-term memory in children with sickle cell disease: executive versus modality-specific processing deficits. *Arch Clin Neuropsychol* 2005; 20: 1073-1085.
60. Jerrell JM, Tripathi A, Stallworth JR. Pain management in children and adolescents with sickle cell disease. *Am J Hematol* 2011; 86: 82-84.
61. Jackson KC, St Onge EL. Antidepressant pharmacotherapy: considerations for the pain clinician. *Pain Pract* 2003; 3: 135-143.
62. Sharp J, Keefe B. Psychiatry in chronic pain: a review and update. *Curr Psychiatry Rep* 2005; 7: 213-219.
63. Barakat LP, Smith-Whitley K, Schulman S, Rosenberg D, Puri R, Ohene-Frempong K. Nocturnal enuresis in pediatric sickle cell disease. *J Dev Behav Pediatr* 2001; 22: 300-305.